

THE COMPLEXITY OF THE CORE MODEL

WILLIAM J. MITCHELL

2/1/2008

ABSTRACT. We use the Σ_3^1 absoluteness theorem to show that complexity of the statement “ (ω, E) is isomorphic to an initial segment of the core model” is Π_4^1 , and the complexity of the statement “ (ω, E) is isomorphic to a member of K ” is Δ_5^1 .

— Complexity of “ x is a real of K ” is Π_3^1 . But we want to look at sets of K which are countable in V .

Definition 1. If Γ is any point class over the reals then we extend the usage of Γ by saying that if $X \subset H_{\omega_1}$ is in Γ if it is each member of X is transitive and $\{ E \subset \omega^2 : (\exists x \in X)(\omega, E) \cong (x, \in) \}$ is in Γ .

Suppose that $K = L[\mathcal{E}]$, so that the initial segments of K are the sets of the form $J_\alpha[\mathcal{E}]$.

Proposition 2.

- (1) “ $\exists \alpha$ “ $x = J_\alpha[\mathcal{E}]$ ” is Π_4^1 .”
- (2) “ x is in K ” is Δ_5^1 .

Proof. We consider clause (1) first. The assertion that x is a premouse, that is, that there is an ordinal α and a good sequence $\mathcal{F}$ of extenders such that $x = J_\alpha(\mathcal{F})$, is Σ_2^1 ; and the assertion that $J_\alpha(\mathcal{F})$ is iterable is Π_2^1 , so the critical element in the characterization of initial segments of K is assertion that $\mathcal{F}$ is maximal. The assertion that $\mathcal{F}$ is maximal for partial extenders, that is, that it contains every mouse which it could have, is “every mouse $J_\nu[\mathcal{G}]$ with projectum ρ such that $\mathcal{G} \restriction \rho = \mathcal{F} \restriction \rho$ is in $J_\alpha[\mathcal{F}]$,” which is Π_3^1 since “ $J_\nu[\mathcal{G}]$ is a mouse” is Π_2^1 .

Thus we need only consider the assertion that $\mathcal{F}$ contains all of the full measures which it should have. By [mit.coreii], this means that $\mathcal{F}$ contains every extender F satisfying

- (1) F could be added to $\mathcal{F} \restriction \gamma$ as $\mathcal{E}_\gamma$, that is, that $\mathcal{F} \restriction \gamma \hat{\cup} F$ is good at γ , and
- (2) If $J_\nu[\mathcal{G}]$ is any iterable premouse such that $\mathcal{G} \restriction \gamma = \mathcal{E} \restriction \gamma$ then $\text{ult}(J_\nu[\mathcal{G}], F)$ is well founded.

The assertion that an extender F satisfies clause (1) is Σ_1^1 . The assertion that a model $J_\alpha[\mathcal{G}]$ is iterable is Π_2^1 , so the assertion that an F satisfies clause (2) is Π_3^1 and hence the assertion that every extender F satisfying clauses (1) and (2) is in $\mathcal{F}$ is Π_4^1 .

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

Since a set $x \in H_{\omega_1}$ is a member of K if and only if x is a member of some countable initial segment of K , it is easy to express “ $x \in K$ ” by a Σ_5^1 formula. To see that “ $x \in K$ ” can also be expressed by a Π_5^1 formula, note that if ν is the rank of x (that is, the rank of the transitive closure of x regarded as a well founded relation) then x is in K if and only if it is in some mouse $J_\alpha[\mathcal{G}]$ such $\mathcal{G} \restriction \nu = \mathcal{E} \restriction \nu$. Thus $x \in K$ if and only if

$$(*) \quad \forall y \forall \nu ((\nu = \text{rank}(x) \text{ and } y = \mathcal{E} \restriction \nu) \implies \exists \mathcal{M} (\mathcal{M} = J_\alpha[\mathcal{G}] \text{ is an iterable mouse with } \mathcal{G} \restriction \nu = y)).$$

By clause (1) of the proposition, $y = \mathcal{E} \restriction \nu$ is Π_4^1 and hence $(*)$ is Σ_5^1 . $\square$

The main result of this paper is that the calculations in the last proposition are best possible. We use the Σ_3^1 -absoluteness theorem from [mit.abs]:

Theorem 3. *Suppose that there is no inner model of $\exists \kappa o(\kappa) = \kappa^{++}$ and that $a^\sharp$ exists for each real a . Then any model M such that $K^{(M)}$ is an iterated ultrapower of K is Σ_3^1 correct.*

Theorem 4.

- (1) *Suppose that M is a model of set theory which satisfies that κ is measurable and that the sharp exists for every subset of κ^+ . Then there is a model in which “ x is an initial segment of K ” is not expressible by a Σ_4^1 formula.*
- (2) *Suppose that M is a model of set theory which satisfies that there are infinitely many measurable cardinals below λ and the sharp exists for every subset of λ . Then there is a model in which “ $x \in K$ ” cannot be expressed by a boolean combination of Σ_4 formulas.*

Proof. Let M satisfy the conditions of clause (1). By taking a submodel if necessary we can assume that $M \models (V = K \text{ and } \neg \exists \kappa o(\kappa) = \kappa^{++})$. Let κ be the smallest measurable cardinal in M , let U be the measure on κ in M , and let $i : M \rightarrow \text{ult}(M, U)$ be the canonical embedding. Finally let λ be the first fixed point of i above κ , so that $\kappa^+ < \lambda < \kappa^{++}$, let $\mathbb{P}$ be the Levy collapse of λ , and let G be $\mathbb{P}$ -generic over M . We will show that there is no Σ_4^1 formula ϕ such that

$$(*) \quad M[G] \models \forall x (\phi(x) \iff x \text{ is an initial segment of } K).$$

Suppose to the contrary that ϕ is such a formula. Since ϕ doesn’t contain any parameters, the homogeneity of the Levy collapse implies that $(*)$ is forced by the empty condition of $\mathbb{P}$. Now $\mathbb{P}$ is also the Levy collapse of $\lambda = i(\lambda)$ in $M_1 = \text{ult}(M, U)$, and G is $\mathbb{P}$ -generic over M_1 , so $(*)$ also holds in $M_1[G]$.

Notice that $M_1[G]$ is a definable submodel of $M[G]$ and $K^{(M_1)} = \text{ult}(M, U)$ is an iterated ultrapower of $M = K^{(M)}$. Furthermore, since M satisfies that every subset of λ has a sharp, it follows that in $M[G]$ every real has a sharp. Thus theorem 3, applied in $M[G]$, implies that every Σ_4^1 formula true in M_1 is true in $M[G]$.

Now let $x = J_\lambda[\mathcal{E}^{M_1}]$, so x is countable in the models M and M_1 . Since x is an initial segment of $K^{(M_1[G])}$ but not of $K^{(M[G])}$ we know that $M_1[G] \models \phi(x)$, while $M \models \neg \phi(x)$, but since ϕ is Σ_4^1 this contradicts the last paragraph, and this contradiction completes the proof of clause (1).

In order to prove clause (2) suppose that M is a model of set theory containing infinitely many measurable cardinals, and that if λ is the sup of the first ω measurable cardinals in M then $x^\sharp \in M$ for each $x \subset \lambda$ in M . As before, we can assume that $M \models (V = K + \neg \exists \kappa o(\kappa) = \kappa^{++})$, so that $M = L[\mathcal{E}]$. Let $\mathbb{P}$ be the partial order to collapse λ , let G be $\mathbb{P}$ -generic over M , and let a be a real which is Cohen generic over $M[G]$. Our model will be $M_0 = M[G, a]$. We will show first that $x \in K$ is not expressible in M_0 by any formula of the form $\pi(x) \wedge \sigma(x)$ where $\pi(x)$ is a Π_4^1 formula and σ is a Σ_4^1 -formula, and afterward we will generalize this to arbitrary boolean combinations of Σ_4^1 formulas.

Suppose then, that $x \in K$ is expressible in M_0 by such a formula $\pi(x) \wedge \sigma(x)$. Let $(U_i : i \in \omega)$ be the first ω measures in $M = L[\mathcal{E}]$, let $i_1 : L[\mathcal{E}] \rightarrow L[\mathcal{E}_1]$ be the iterated ultrapower using each of the measures U_i exactly once, let $i_a : L[\mathcal{E}] \rightarrow L[\mathcal{E}_a]$ be the iterated ultrapower using only the measures U_i for $i \in a$, and Set $N = L[\mathcal{E}_a][G, a]$ and $M_1 = L[\mathcal{E}_1][G, a]$. Then

$$M_0 \supset N \supset M_1.$$

Then $\mathcal{E}_1 \restriction \lambda$ is a member of $K^{(M_0)}$ and $K^{(M_1)}$, but not a member of $K^{(N)}$. Just as in the proof of clause (1), the formula $(**)$ holds in N and M_1 , so

$$(***) \quad \pi(\mathcal{E}_1 \restriction \lambda) \wedge \sigma(\mathcal{E}_1 \restriction \lambda)$$

is true in M_0 and M_1 but false in N . But lemma 3 implies that M_1 is Σ_3^1 correct in N , so that

$$M_1 \models \sigma(\mathcal{E}_1 \restriction \lambda) \implies N \models \sigma(\mathcal{E}_1 \restriction \lambda)$$

and N is Σ_3^1 correct in N so

$$M_1 \models \pi(\mathcal{E}_1 \restriction \lambda) \implies N \models \pi(\mathcal{E}_1 \restriction \lambda),$$

contradicting formula $(***)$ and hence disproving $(**)$.

Now suppose that

$$M_0 \models \forall x (x \in K \iff \phi(x))$$

where $\phi(x)$ is a boolean combination of Σ_4^1 formulas. Then $\phi(x)$ is equivalent to a formula of the form

$$\bigvee_{i < n} (\pi_i(x) \wedge \sigma_i(x))$$

for some $n \in \omega$ and formulas $\pi_i(x)$ and $\sigma_i(x)$ which are Π_3^1 and Σ_3^1 , respectively. Now we define a chain

$$M_0 \subset N_0 \subset M_1 \subset \cdots \subset N_{n-1} \subset M_n$$

of models. As before, we have $M_0 = M[G, a] = L[\mathcal{E}][G, a]$. Let $j_i : L[\mathcal{E}] \rightarrow L[\mathcal{E}_i]$ be the iterated ultrapower using each of the measures $(U_i : i \in \omega)$ exactly i times, and let $L[\mathcal{E}_{i,a}]$ be the result of iterating each of the measures $\{j_i(U_k) : k \in a\}$ once more. As before, $\mathcal{E}_n$ will be a member of M_i for each $i \leq n$, and formula (4) is true for each of the models M_i so that $\phi(\mathcal{E}_i \restriction \lambda)$ is true in each M_i . Then by the pigeon hole principle there are integers $k < n$ and $i_0 < i_1 \leq n$ such that $\sigma_k(\mathcal{E}_i \restriction \lambda) \wedge \pi_k(\mathcal{E}_i \restriction \lambda)$ is true in both the models M_{i_0} and M_{i_1} . Just as in the last paragraph this implies

that $\sigma_k(\mathcal{E}_i \restriction \lambda) \wedge \pi_k(\mathcal{E}_i \restriction \lambda)$ is true in N_{i_0} , but this is impossible because $\mathcal{E}_n$ is not in $K^{(N_{i_0})}$. $\square$

The hypothesis cannot be weakened in theorem 4. Taking clause (1) for example, if N is any model such that $K^{(N)}$ doesn't have any measurable cardinals which are countable in N then x is an initial segment of $K^{(N)}$ if it contains all the mice it should, which is a Π_3^1 statement; while if $K^{(N)} = L[\mathcal{E}]$ where $\mathcal{E}$ is countable in N then $\mathcal{E}$ is characterized by the fact that $\mathcal{E}^\sharp$ doesn't exist and $\text{ult}(L_{\omega_1}(\mathcal{E}), E)$ is ill founded for any ultrafilter E on $L[\mathcal{E}]$ which is not either on the sequence $\mathcal{E}$ or an iterate of a measure which is on $\mathcal{E}$. The hypothesis to clause (1) is the weakest which will allow the existence of a model N such that $K^{(N)}$ contains a countable mouse and is not equal $L[\mathcal{E}]$ for a sequence $\mathcal{E}$ which is countable in N .

Similarly, the boldface version of theorem 4 requires that the measurable cardinals of $K^{(N)}$ be cofinal in $\omega_1^{(N)}$, which requires that we start with an inaccessible limit of measurable cardinals. The arguments above readily yield

Theorem 5. *Suppose that there is a model M of set theory with an inaccessible limit of measurable cardinals. Then there is a model N such that “ x is an initial segment of K ” is not expressible in N by a Σ_4^1 formula, and “ $x \in K$ ” cannot be expressed in N by any boolean combination of Σ_4 formulas.*